



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY  
AND POLLUTION PREVENTION

August 10, 2015

Nathan P. Ehresman  
Director, Regulatory Affairs  
Nufarm Americas Inc – AGT Division  
c/o Nufarm Americas Inc.  
4020 Aerial Center Parkway  
Morrisville, NC 27560

Subject: Label Amendment – Revision of Crop Use Listing (update Seed & Pod Vegetable to Legume Vegetable Crop Group)  
Product Name: Sebring 480 FS Fungicide  
EPA Registration Number: 55146-106  
Application Dated: 05/04/2015  
Decision Number: 505277

Dear Mr. Ehresman:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. “To distribute or sell” is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

Should you wish to add/retain a reference to the company’s website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product’s label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA’s Office of Enforcement and Compliance.

Page 2 of 2  
EPA Reg. No. 55146-106  
Decision No. 505277

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. If you have any questions, please contact Marcel Howard by phone at (703)305-6784, or via email at [howard.marcel@epa.gov](mailto:howard.marcel@epa.gov).

Sincerely,

A handwritten signature in black ink, appearing to be 'HJ', written in a cursive style.

Hope Johnson, Product Manager 21  
Fungicide Branch  
Registration Division (7505P)  
Office of Pesticide Programs

Enclosure: Stamped "Accepted" Product Labeling.

# SEBRING™ 480 FS FUNGICIDE

FOR USE AS A SEED TREATMENT FOR CONTROL OF SYSTEMIC DOWNY MILDEW,  
*PYTHIUM* SEED ROT, *PYTHIUM* DAMPING-OFF, AND EARLY SEASON *PHYTOPHTHORA*  
DISEASES OF CERTAIN CROPS.

**ACTIVE INGREDIENT:**

Metalaxyl: N-(2,6-dimethylphenyl)-N-(methoxyacetyl) alanine methyl ester ..... 44.08%

**OTHER INGREDIENTS:** ..... 55.92%

**TOTAL:** ..... 100.00%

Contains 4.0 pounds Metalaxyl per gallon (480 grams per liter).

KEEP OUT OF REACH OF CHILDREN

## CAUTION

SEE INSIDE BOOKLET FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300  
For Medical Emergencies Only, Call (877) 325-1840

**A C C E P T E D**

08/10/2015

Under the Federal Insecticide, Fungicide  
and Rodenticide Act as amended, for the  
pesticide registered under  
EPA Reg. No. 55146-106

EPA REG. NO. 55146-106  
EPA EST. NO. \_\_\_\_\_

Manufactured For  
NUFARM AMERICAS INC.  
AGT Division  
11901 S. Austin Avenue  
Alsip, IL 60803



**NET CONTENTS:** \_\_\_\_\_ GAL. (\_\_\_\_\_ Liters)  
[Designation as "NONREFILLABLE" or "REFILLABLE" for containers > 5 GAL]

<b>FIRST AID</b>	
<b>IF SWALLOWED</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• <b>DO NOT</b> induce vomiting unless told to by the poison control center or doctor.</li> <li>• <b>DO NOT</b> give anything to an unconscious person.</li> </ul>
<b>IF IN EYES</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF ON SKIN OR CLOTHING</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15 to 20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF INHALED</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth- to-mouth, if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>
<b>HOT LINE NUMBER</b>	
<p>Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.</p>	

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS  
CAUTION**

Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

**PERSONAL PROTECTIVE EQUIPMENT (PPE)**

**Applicators and other handlers must wear:**

- Long-sleeved shirt and long pants,
- Shoes and socks,
- Chemical-resistant gloves: barrier laminate or nitrile rubber ≥ 14 mils or neoprene rubber ≥ 14 mils or Viton ≥ 14 mils.

**User Safety Requirements:**

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

<b>USER SAFETY RECOMMENDATIONS</b>
<p><b>Users Should:</b></p> <ul style="list-style-type: none"> <li>• Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.</li> <li>• Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.</li> <li>• Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.</li> </ul>

## ENVIRONMENTAL HAZARDS

**DO NOT** apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. **DO NOT** apply when weather conditions favor drift from treated areas. **DO NOT** contaminate water when disposing of equipment washwater or rinsate.

Groundwater Advisory Statement: This chemical is known to leach through soil into ground water under certain conditions as a result of agricultural use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination.

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**DO NOT** apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

## AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

**DO NOT** enter or allow worker entry into treated areas during the **restricted-entry interval (REI) of 24 hours**.

Exception: If the seed is treated with the product and the treated seed is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water is: coveralls, shoes plus socks and chemical-resistant gloves: barrier laminate or nitrile rubber  $\geq 14$  mils or neoprene rubber  $\geq 14$  mils or Viton  $\geq 14$  mils.

## PRODUCT INFORMATION

This product is a systemic fungicide seed dressing specifically for control of systemic downy mildews, *Pythium* and *Phytophthora spp.* For control of other soilborne diseases, combination of captan, thiram, and carboxin registered formulations are compatible. **DO NOT** use with other seed treatment products unless previous experience assures compatibility.

### Precautions:

**Note:** This product is a systemic fungicide having a specific mode of action. This product could be subject to development of resistant strains of fungi. Development of resistance cannot be predicted. Consult with your State Agricultural Experiment Station or Extension Service Specialist for guidance and ways to control any possible metalaxyl resistant strains of fungi which may occur. **FAILURE TO FOLLOW DIRECTIONS OF THIS LABEL MAY RESULT IN CROP INJURY OR POOR DISEASE CONTROL.**

### Restrictions:

**In California:** Only for use in commercial seed treating establishments. Not allowed for "on-farm" use.

**On-Farm Seed Treatment: DO NOT** apply this product through any type of irrigation system.

**DO NOT** use treated seed for food or feed purposes or process for oil. Treat only those seeds needed for immediate use, minimizing the interval between treatment and planting. **DO NOT** store excess treated seeds beyond planting time.

Seed treatment on agricultural establishment at or immediately before planting is within the scope of WPS, while commercial treatment of seeds is not within the scope.

**Planter Box Treatment:** Apply at the specified rate and premix with the seed directly in the planter box at planting. Place half the amount of seed into the planter box and apply half the required amount of pesticide to the seed. Stir the seed thoroughly with a paddle until the seed is uniformly coated. Place the remaining half of the seed into the planter box and add the remaining half of the pesticide. Again stir the seed with the paddle until all seed is thoroughly coated.

**For Commercial Seed Treatment:** Seed treated commercially or for sale must be dyed an unnatural color using an EPA approved dye before sale. Federal law requires that bags containing treated seed shall be labeled with the following information:

**Tank Mixing:** When tank mixing follow the most restrictive label instructions for any tank mix partner. **DO NOT** tank mix with any product that contains a prohibition against tank mixing.

### **SEED BAG LABELING REQUIREMENTS**

Seed commercially treated with this product must be labeled in accordance with all applicable requirements of the Federal Seed Act. The user of this product is responsible for ensuring that the seed bag meets all requirements under the Federal Seed Act.

THE FEDERAL SEED ACT REQUIRES THAT BAGS CONTAINING TREATED SEEDS BE LABELED WITH THE FOLLOWING STATEMENTS:

*This seed has been treated with **Sebring 480 FS Fungicide** (metalaxyl).*

**DO NOT** use for feed, food, or oil purposes.

THE US ENVIRONMENTAL PROTECTION AGENCY REQUIRES THE FOLLOWING STATEMENTS ON BAGS CONTAINING SEEDS TREATED WITH **Sebring 480 FS Fungicide** (metalaxyl):

*Store treated seed away from feed or foodstuffs.*

**DO NOT** allow children, pets, or livestock to have access to treated seeds.

**DO NOT** use the same auger for treating seed that will handle commercial or feed grains.

**DO NOT** plant any crop not listed on this label within 30 days of planting **Sebring 480 FS Fungicide**-treated seed.

**DO NOT** harvest mustard greens. Rape greens grown and harvested from **Sebring 480 FS Fungicide**-treated seed must not be used for human and feed consumption. Rapeseed grown and harvested from **Sebring 480 FS Fungicide**-treated seed is only for industrial uses and cannot be used for edible oil or any other human/feed consumption.

**DO NOT** use Metalaxyl or mefenoxam for control of Pythium in water-seeded rice.

*Wear long pants, long-sleeved shirt, shoes plus socks, and chemical-resistant gloves made of any waterproof material when opening this bag or handling (e.g., loading, pouring) treated seed.*

*After treated seed has been planted, **DO NOT** enter or allow worker entry into areas planted with treated seeds during the restricted-entry interval (REI) of 24 hours. **EXCEPTION:** Workers may enter these areas without restriction if there will be no worker contact with the soil/media subsurface.*

Treated seed exposed on soil surfaces may be hazardous to wildlife. Cover or collect treated seed spilled during loading and planting.

**DO NOT** contaminate bodies of water when disposing of planting equipment washwater. If excess treated seed remains after planting, dispose of it by burying away from bodies of water.

**DO NOT** store leftover treated seed where there is a chance of it becoming mixed with untreated seed.

Store treated seed in cool, dry conditions.

Dispose of treated seed packaging in accordance with local requirements.

Excess treated seed may be used for ethanol production only if (1) by-products are not used for livestock feed and (2) no measurable residues of pesticides remain in the ethanol by-products that are used in agronomic practice.

### SEED TREATMENT

This product may be applied as a water-based slurry with other registered seed treatment insecticides and fungicides through standard slurry or mist-type commercial seed treatment equipment.

### DOMESTIC USE

CROP	DISEASE	RATE (Fluid Ounces per 100 Pounds of Seed)
<b>Brassica Leafy Vegetables</b> Broccoli (including Chinese and raab), Brussels sprout, Cabbage (including Chinese bok choy, Chinese napa and mustard), Cauliflower, Cavalo broccoli, Collards, Kale, Kohlrabi, Mizuna, Mustard greens, Mustard Spinach, and Rape greens)	<i>Pythium</i> damping-off control	0.50
<b>Beets, Garden &amp; Sugarbeets</b>	<i>Pythium</i> damping-off control	0.50
<b>Canola*</b> <b>Mustard Seed*</b> <b>Rapeseed*</b>	<i>Pythium</i> damping-off control	0.20 - 0.30
	<b>Restriction: DO NOT</b> harvest mustard greens. Rape greens grown and harvested from Sebring 480 FS Fungicide treated seed must not be used for human and feed consumption. Rapeseed grown and harvested from Sebring 480 FS Fungicide treated seed is only for industrial uses and cannot be used for edible oil or any other human/feed consumption.	
<b>Carrots*</b>	<i>Pythium</i> damping-off control	0.50
<b>Cotton</b>	<i>Pythium</i> seed rot and damping-off control	0.50 - 1.00

CROP	DISEASE	RATE (Fluid Ounces per 100 Pounds of Seed)	
<b>Cucurbit Vegetables</b> Chayote (fruit), Chinese waxgourd (Chinese preserving melon), Citron melon, Cucumber, Gherkin, Gourd (edible) (includes hyotan, cucuzza, hechima, Chinese okra), <i>Momordica</i> spp (includes balsam apple, balsam pear, bittermelon, Chinese cucumber), Muskmelon (including hybrids and/or varieties of <i>Cucumis melo</i> (including true cantaloupe, cantaloupe, casaba, Santa Claus melon, crenshaw melon, honeydew melon, honey balls, Persian melon, golden pershaw melon, mango melon, pineapple melon, snake melon), Watermelon (including hybrids and/or varieties of <i>Citrullus</i> spp.), Pumpkin, Squash (summer, winter including butternut squash, calabaza, hubbard squash, acorn squash, spaghetti squash).	<i>Pythium</i> damping-off control	0.50	
<b>Forage Grasses</b> Grasses grown for hay, grazing or silage, Corn (fodder or silage), Sorghum (hay or silage), and Small grain grown for hay, grazing or silage.	<i>Pythium</i> damping-off control	0.50	
<b>Forage Legumes</b> Alfalfa, Beans (for forage), Clover, Cowpeas (for forage), Cowpea hay, Lespedeza, Peanuts (for forage), Peanut hay, Peas (for forage), Pea vine hay, Soybeans (for forage), Soybean hay, Trefoil, Velvet beans (for forage) and Vetch.	<i>Pythium</i> damping-off and early season <i>Phytophthora</i> control	0.50 - 1.00	
<b>Fruiting Vegetables</b> Eggplant, Ground cherry ( <i>Physalis</i> spp), Pepino, Pepper (includes bell pepper, chili pepper, cooking pepper, pimento, sweet pepper), Tomatillo, Tomato.	<i>Pythium</i> damping-off control	0.50	
<b>Grain Crops</b> Barley, Buckwheat, Corn (field corn, sweet corn and popcorn), Millet, Milo, Oats, Rye, Triticale, and Wheat.	<i>Pythium</i> damping-off control	0.50	
	Systemic downy mildew control of sweet corn	1.00	
<b>Use Rate Conversions for Field Corn</b>			
<b>PRODUCT FLUID OUNCES /</b>			
<b>100 lb seed<sup>†</sup></b>	<b>1000 seeds</b>	<b>80,000 seed count</b>	<b>MG AI / SEED</b>
0.50	0.0028	0.224	0.040
1.00	0.0056	0.448	0.080
This product contains 14,195.5 mg metalaxyl (AI) / fluid ounce. <sup>†</sup> Based on standard number of 1786 seeds / lb (100,000 seeds / 56 lb bu)			
<b>NOTE:</b> To maintain mg ai / seed, adjust fluid ounces / 100 lb in proportion to the actual seeds / lb compared to the standard number of seeds / lb for that crop. For example if there are 1600 corn seeds / lb and the targeted rate is 0.040 mg ai / seed at a non adjusted rate of 0.50 fl. oz / 100 lb. of seed, use: $(1600 \div 1786) \times 0.50 = 0.45$ fl. oz / 100 lb of seed.			
<b>Leafy Vegetables</b>	<i>Pythium</i> damping-off control	0.50	

CROP	DISEASE	RATE (Fluid Ounces per 100 Pounds of Seed)
<p>Amaranth (Chinese spinach), Cardoon, Celery, Celery (Chinese), Celtuce, Chervil, Chrysanthemum (edible-leaved, garland), Corn salad, Cress (garden, upland), Dandelion, Dock (sorrel), Endive (escarole), Fennel (Florence), Lettuce (head and leaf), Orach, Parsley, Purslane (garden, winter), Radicchio (red chicory), Rhubarb, Spinach (including New Zealand and vine), Swiss chard.</p>	<p><b>Restriction: DO NOT</b> apply this product to spinach which is destined to be grown in the greenhouse. Containers or bags containing Sebring 480 FS treated spinach seed must be labeled with this restriction.</p>	
<p><b>Legume Vegetables (except soybeans)</b>            Bean (<i>Lupinus</i> spp.) [includes grain lupin, sweet lupin, white lupin, white sweet lupin]; Bean (<i>Phaseolus</i> spp.) [includes field bean, kidney bean, lima bean (succulent or dry), navy bean, pinto bean, runner bean, snap bean, tepary bean, wax bean]; Bean (<i>Vigna</i> spp.) [includes adzuki bean, asparagus bean, blackeyed pea (succulent or dry), catjang, Chinese longbean, cowpea (succulent or dry), Crowder pea, moth bean (succulent or dry), mung bean, rice bean, southern pea (succulent or dry), urd bean, yardlong bean]; Broad bean (fava) (succulent or dry); Chickpea (garbanzo bean); Guar; Jackbean; Lablab bean (hyacinth); Lentil; Pea (<i>Pisum</i> spp.) [includes dwarf pea, edible-pod pea, English pea, field pea, garden pea, green pea, snow pea, sugar snap pea]; Pigeon pea (succulent or dry); Sword bean.</p>	<p><i>Pythium</i> damping-off control</p>	<p>0.50</p>
	<p>Early season <i>Phytophthora</i> control</p>	<p>0.50</p>
	<p>Control of <i>Pythium</i> damping-off and systemic downy mildew of peas</p>	<p>1.00</p>
<p><b>Onions (dry bulb and green)</b></p>	<p><i>Pythium</i> damping-off control</p>	<p>0.50</p>
<p><b>Peanuts*</b></p>	<p><i>Pythium</i> seed rot and damping-off control</p>	<p>0.50</p>
<p><b>Rice*</b></p>	<p><i>Pythium</i> damping-off control</p>	<p>0.50 - 1.00</p>
	<p><b>Restriction:</b> If using Sebring 480 FS treated rice do not use metalaxyl or mefenoxam for control of <i>Pythium</i> in water-seeded rice. Containers or bags containing Sebring 480 FS treated rice seed must be labeled with this restriction.</p>	
<p><b>Root and Tuber Vegetables (except potato)</b>            Arracacha, Arrowroot, Artichoke (Chinese, Jerusalem), Beet (garden, sugar), Burdock (edible), Canna (edible), Carrot, Cassava (bitter and sweet), Celeriac, Chayote (root), Chervil (turnip-rooted), Chicory, Chufa, Dasheen (taro), Ginger, Ginseng, Horseradish, Leren, Parsley (turnip-rooted), Parsnip, Radish (including oriental), Rutabaga, Salsify (including black and Spanish), Skirret, Sweet potato, Tanier, Turmeric, Turnip, Yam bean, Yam (true).</p>	<p><i>Pythium</i> damping-off control</p>	<p>0.50</p>

CROP	DISEASE	RATE (Fluid Ounces per 100 Pounds of Seed)	
Sorghum	<i>Pythium</i> damping-off control	0.25 - 0.50	
	Systemic downy mildew (races susceptible to chemical control)	0.50 - 2.00	
	<b>Note:</b> Apply 0.50 – 1.00 fluid ounces of this product on cultivars which possess sorghum downy mildew resistance. To those cultivars which do not possess resistance to a particular race of sorghum downy mildew in heavily infected areas with recent severe history, apply 1.00 – 2.00 fluid ounces of this product. Only high quality sorghum seed (90% germination and above) should be treated with this product.		
Soybeans	<i>Pythium</i> damping-off control and early season <i>Phytophthora</i> control	0.50 - 1.00	
<b>Use Rate Conversions for Soybeans</b>			
<b>PRODUCT FLUID OUNCES /</b>			
<b>100 lb seed<sup>†</sup></b>	<b>1000 seeds</b>	<b>80,000 seed count</b>	<b>MG AI / SEED</b>
0.50	0.0017	0.136	0.024
1.00	0.0034	0.272	0.048
This product contains 14,195.5 mg metalaxyl (AI) / fluid ounce. <sup>†</sup> Based on standard number of 3000 seeds / lb (150,000 seeds / 50 lb bu) <b>NOTE:</b> To maintain mg ai / seed, adjust fluid ounces / 100 lb in proportion to the actual seeds / lb compared to the standard number of seeds / lb for that crop. For example if there are 2800 soybean seeds / lb and the targeted rate is 0.024 mg ai / seed at a non adjusted rate of 0.50 fl. oz / 100 lb. of seed, use: $(2800 \div 3000) \times 0.50 = 0.467$ fl. oz / 100 lb of seed.			
Spinach*	<i>Pythium</i> damping-off control and early season <i>Phytophthora</i> control	0.50 – 1.00	
	<b>Restriction: DO NOT</b> apply this product to spinach which is destined to be grown in the greenhouse. Containers or bags containing Sebring 480 FS treated spinach seed must be labeled with this restriction.		
Sunflowers	Systemic downy mildew control	1.00 - 2.00	
<b>Turf Grasses</b> Golf Courses, Home Lawns, Reclamation and Erosion Control Projects	<i>Pythium</i> damping-off control	1.00	

\* Not registered for use in California

### FOR EXPORT USE ONLY

For treatment of the following seed for export to countries requiring treatment with metalaxyl before importation.

CROP	DISEASE	RATE (Fluid Ounces per 100 Pounds of Seed)
<b>Brassica (Cole) Leafy Vegetables</b>	<i>Pythium</i> damping-off control	0.50 - 1.00
<b>Canola</b>	<i>Pythium</i> damping-off control	0.30 - 1.10 (20 - 66 ml per 100 Kg of seed)
<b>Corn</b>	<i>Pythium</i> damping-off control	0.50 - 1.00
	Systemic downy mildew control	2.00
<b>Cucurbit Vegetables</b>	<i>Pythium</i> damping-off control	0.50
<b>Fruiting Vegetables</b>	<i>Pythium</i> damping-off control	0.50 - 1.00
<b>Leafy Vegetables</b>	<i>Pythium</i> damping-off control	0.50 - 1.00
<b>Millet</b>	<i>Pythium</i> damping-off control	0.50 - 1.00
	Systemic downy mildew control	1.70
<b>Onions</b>	<i>Pythium</i> damping-off control	0.50 - 1.00
<b>Peas</b>	<i>Pythium</i> damping-off control and early season <i>Phytophthora</i> control	2.30
<b>Root and Tuber Vegetables (except potato)</b>	<i>Pythium</i> damping-off control	0.50 - 1.00
	<b>Restriction: DO NOT</b> apply this product to potatoes.	
<b>Sorghum</b>	<i>Pythium</i> damping-off control	0.50 - 1.00
	Systemic downy mildew control	3.50
<b>Sunflower</b>	Systemic downy mildew control	6.00

Seed treated in accordance with the export use pattern must be labeled as follows:

**FOR EXPORT ONLY  
NOT FOR DOMESTIC SALES OR USE**

**DIRECTIONS FOR USE AT REDUCED RATES IN COMBINATION WITH OTHER FUNGICIDES**

This product is a systemic fungicide seed dressing which may be applied in combination with other EPA-registered fungicides to aid in control of *Pythium*. This product may be applied as a water-based slurry through standard slurry of mist-type commercial seed treatment equipment.

CROP	DISEASE	RATE (Fluid Ounces per 100 Pounds of Seed)
<b>Barley, Corn (field), Oats, Rye and Wheat</b>	Aid in control of seed decay and damping-off caused by <i>Pythium</i>	0.07- 0.25
<b>Beans (field, kidney, navy and pinto), Lentils, Peas and Soybeans</b>		
<b>Peanuts*</b>		
<b>Rice*</b>		
<b>Sunflowers</b>		

\* Not registered for use in California

**NOTE:** Apply only in combination with EPA registered rates of broad-spectrum seed treatment fungicides.

## STORAGE AND DISPOSAL

**DO NOT** contaminate water, food or feed by storage or disposal.

**PESTICIDE STORAGE:** Store in cool dry location. **DO NOT** store in direct hot sunlight. **DO NOT** store above 90°F for prolonged periods of time. After prolonged storage in intense cold, allow product to return to approximately 75°F before using.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

### CONTAINER HANDLING:

**[Note to Reviewer:** The following statement will be included on all Final Printed Labels bearing multiple Container Disposal (Container Handling) statements] **NOTE:** This product is available in multiple containers. Refer to the Net Contents section of this products labeling for the applicable “Nonrefillable” or “Refillable” designation. Follow the container disposal [handling] instructions below that apply to your container type / size.”

#### [Nonrefillable Containers 5 Gallons or Less]

**Nonrefillable container. DO NOT** reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying.

**Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke.

#### [Nonrefillable containers larger than 5 gallons]

**Nonrefillable container. DO NOT** reuse or refill this container. Offer for recycling if available. Triple rinse or pressure rinse container (or equivalent) promptly after emptying.

**Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke.

**Pressure rinse as follows:** Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

#### [Refillable containers]

**Refillable container.** Refill this container with pesticide only. **DO NOT** reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. Agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

#### [Refillable For return to Nufarm:]

**Refillable Container.** Refill this container with pesticide only. **DO NOT** reuse this container for any other purpose. Close all openings and replace all caps. Contact Nufarm’s Customer Service Department at 1-800-345-3330 to arrange for return of the empty refillable container.

## WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

## LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR ARISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

**If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.**

(RV080515)

Sebring is a trademark of Nufarm Americas Inc.

## LABEL HISTORY

<u>File Name</u>	<u>Revision Mark</u>	<u>Comment</u>
055146-00106.20120427.Revised_Label	RV042712	Addition of Use Rate conversions for Corn & Soybeans Update of Container Disposal Statement to add flexibility for single label on multiple container types
055146-00106.20120802.MASTER	RV080212	APPROVAL
055146-00106.20120827.Notification	RV082712	In Domestic Use table, remove California restriction on Brassica Leafy Vegetables, Cucurbits, Leafy Vegetables, Fruiting Vegetables, Onions, and Root and Tuber Vegetables (except potatoes). Change * to † rate table to avoid confusion with asterisk.
055146-00106.20120827b.Notification	RV082712	10-03-12: per EPA changed to Amendment. Move Celeriac from Leafy Vegetables table to Root and Tuber Vegetables table.
055146-00106.20121010.MASTER	RV101012	EPA Approval
055146-00106.20150504.DRAFT	RV050415	Draft Label- revised to Seed and Pod Vegetables to Legume Vegetables Crop Group
055146-00106.20150805.DRAFT	RV080515	Revisions made Per EPA request 8/4/15